



Bulkamid™

Urethral Bulking System

Ready to say goodbye to bladder leaks?

Say hello to Bulkamid™

Bulkamid is a minimally invasive and long-lasting treatment option for bladder leaks caused by stress urinary incontinence (SUI).

What is stress urinary incontinence (SUI)?

Stress urinary incontinence (SUI), sometimes called stress incontinence, is a condition where unwanted bladder leaks happen during normal daily activities such as coughing, laughing, or exercise.

How can Bulkamid help?

Bulkamid is a urethral bulking agent that is used to treat stress urinary incontinence. It is a soft, water-based gel that can be used to restore the natural closing of the urethra. Similar to facial fillers, Bulkamid adds volume to the surrounding tissue.

Discover the benefits of Bulkamid

Safe and effective

Bulkamid is a minimally-invasive long-lasting treatment option to treat bladder leaks caused by stress urinary incontinence (SUI).²

Long-lasting relief

Designed to effectively treat women suffering from SUI, with clinical support up to 7 years.²


Do you struggle with bladder leaks when you:



Sneeze



Exercise



Laugh




Cough



Walk


You're not alone!



1 in every 2 U.S. women experience some degree of involuntary leakage of urine at some point in their life.¹

Don't settle for frustrating urinary symptoms or incontinence pads.

Ask your doctor today if Bulkamid is right for you.



The procedure

The procedure is a series of 3-4 injections in an outpatient setting under local anesthetic.

To learn more about Bulkamid,
visit www.fixincontinence.com/bulkamid

References:

1. Patel DA, Sutcliffe S, Kelley EP, et al. Updated prevalence of urinary incontinence in women: 2015 -2018. Am J Obstet Gynecol. 2024;230(5):548.e1-548.e10. doi:10.1016/j.ajog.2023.11.005)
2. Brosche T, Kuhn A, Lobodasch K, Sokol ER. Seven-year efficacy and safety outcomes of Bulkamid for the treatment of stress urinary incontinence. Neurourol Urodyn. 2021;40(1):502-508. doi:10.1002/nau.24589

Important Safety Information:

Bulkamid Urethral Bulking System is indicated for the treatment of stress urinary incontinence (SUI) due to intrinsic sphincter deficiency (ISD) in adult women. The Cystoscope is indicated for endoscopic surgical procedures.

Bulkamid Urethral Bulking System must not be used in patients suffering from acute urinary tract infection. Consult your physician if you are suffering from other pelvic infections, pelvic trauma, or pelvic inflammatory disease.

Potential risks may include infection, inability or difficulty to urinate, painful urination, excreted bulking material, embolism, blood in the urine, frequent urination at night, urinary tract infection, intense sudden need to urinate, infection or irritation in the vagina/genital area, worsening urinary incontinence, abnormal blood lab results (Immunoglobulin E), back/neck pain, dizziness/fainting, pain, blockage of blood vessels by a clot, redness of the skin, inflammation, migration of injected Bulkamid material, and urethral erosion.

Patients suffering from blood coagulation disorders are at higher risk of complications including hematoma or urethral bleeding, as with any invasive procedure. If a patient has undergone major surgery or dental work, Bulkamid should not be injected until the patient has fully recovered. The effect of Bulkamid has not been evaluated in women during pregnancy, delivery or lactation. Patients should be counseled that one or more repeat Bulkamid Hydrogel injection procedures may be required.

Be sure to talk with your doctor so that you thoroughly understand all of the risks and benefits associated with the use of the device.

**Boston
Scientific**
Advancing science for life™

Boston Scientific Corporation
300 Boston Scientific Way
Marlborough, MA 01752-1234
www.bostonscientific.com

© 2026 Boston Scientific Corporation
or its affiliates. All rights reserved.

URO-2237808-AB JUN 2026